

LASER IRIDOTOMY

LASER IRIDOTOMY is a procedure performed for prevention of Acute Congestive Glaucoma, which is a sight threatening medical emergency

PREPARATION GUIDE

The following will guide you through each step prior to, during, and after the Iridotomy procedure.

BEFORE YOUR PROCEDURE

- Your Iridotomy appointment is scheduled for _____ at **Samyak Drishti Eye Clinic**.
- You should arrive 30 minutes BEFORE your scheduled appointment time (unless you are otherwise instructed)
- If you must reschedule or cancel your scheduled Iridotomy procedure, please notify us at least 2 days in advance.
- Please arrange to have someone drive you home as your vision may be hazy.

BILLING AND PAYMENTS

- Please make arrangements for payment to Samyak Drishti Eye Clinic prior to the date of your procedure.
- If you are paying by cash, VISA or MasterCard you can pay on the day of surgery, before the procedure
- If you are paying by NEFT, RTGS, Net Banking, electronic fund transfer or online payments, you can pay 12 days before surgery.

The details for the same are:

he details for the same are:

Samyak Drishti Eye Clinic

Bank details: HDFC Bank

Branch: Chowpatty

RTGS / NEFT / IFSC Code: HDFC0001201

A/c No: 50200031445399

□ If you are paying by cheque, you can pay 3 days before the surgery. Cheques can be made in favour of “SAMYAK Drishti Eye Clinic”.

ON THE DAY OF YOUR PROCEDURE

- Have a light meal and continue all medication as advised
- Instil all the eyedrops as advised
- Once you arrive, the staff will assist you with some drops prior to the Iridotomy procedure
- The procedure will take about 10 minutes.
- Your expected time of stay at the Samyak Drishti Eye Clinic will be approximately 1-2 hours.

AFTERCARE

- You will most likely experience some discomfort after this procedure for several hours.
- Your vision may be blurry for 1 to 2 days after your procedure.
- You will be given an anti-inflammatory drop to use 4 times a day for 1 week, including the day of your procedure.

If you have any problems, we can be reached during the day by calling at 022-23613937 / 022-23623937

MEDICATION:

- **Prednisolone Acetate 1% (Pred-Forte):** Anti-inflammatory eye drops which help heal the eye. Apply 1 drop to the operated eye(s) 4 times a day (breakfast, lunch, dinner, and bedtime) for 7 days, including the day of your procedure. **SHAKE WELL** before each use (20 times).
- **Tablet Crocin / Dolo / Lanol-ER (or your preferred pain reliever):** As needed for discomfort. Do not take more than 1 pain reliever at a time.